arXiv:0811.3352v1 [cond-mat.other] 20 Nov 2008

Weizsäcker energy of unitary Fermi gas in a harmonic trap

Alexander L. Zubarev*
Department of Physics, Purdue University
West Lafayette, Indiana 47907

The universal method of construction of the rigorous lower bounds to the Weizsäcker energy is presented. We study a few-fermion system at the unitarity. Upper and lower bounds to the density functional theory (DFT) ground state energy within the local density approximation (LDA) are given. The rigorous lower bounds to the accuracy of the method are derived.

PACS numbers: 31.15.Ew, 71.15.Mb, 03.75.Ss

---

* e-mail: zubareva@physics.purdue.edu

There has been a lot of interest in systems of fermions at the unitarity [1-3] (when the scattering length diverges, the Bertsch many-body problem [4]). While Refs.[5-8] consider homogeneous systems, Refs.[9-13] present *ab initio* calculations of the properties of trapped fermionic atoms.

The modern DFT is based on the Kohn-Sham approach [14], where the noninteracting kinetic energy, T, is calculated in terms of the Kohn-Sham orbitals, although Ref.[15] proved the basic existence of functional $T(\rho)$, where ρ is the density, $\int \rho(\vec{r})d^3r = N$, and N is the particle number. The accurate density-functional approximation to the kinetic energy would reduce dramatically complexity of the DFT calculations (here we note the superfluid extension of the DFT given in Refs.[16-18]). For applications of the DFT to the nuclear structure physics see web site, constructed for the universal nuclear energy density functional (UNEDF) collaboration, http://unedf.org.

The kinetic energy functional cam be written as

$$T[\rho] = \frac{\hbar^2}{2m}\int \tau(\rho(\vec{r}))d^3r, \tag{1}$$

where the semiclassical expansion for the kinetic energy is [19-23]

$$T[\rho] = \frac{\hbar^2}{2m}\int (\tau_{TF}(\rho) + \tau_2(\rho) + \tau_4(\rho) + ...)d^3r, \tag{2}$$

where

$$\tau_{TF}(\rho) = \frac{3}{5}(3\pi^2)^{2/3}\rho^{5/3}, \tag{3}$$

$$\tau_2(\rho) = \frac{1}{36}\frac{(\vec{\nabla}\rho)^2}{\rho}, \tag{4}$$

$$\tau_4(\rho) = \frac{1}{6480}(3\pi^2)^{-2/3}\rho^{1/3}[8(\frac{\vec{\nabla}\rho}{\rho})^4 - 27(\frac{\vec{\nabla}\rho}{\rho})^2\frac{\triangle\rho}{\rho} + 24(\frac{\triangle\rho}{\rho})^2]. \tag{5}$$

The Weizsäcker energy $T_W[\rho]$ [24]

$$T_W[\rho] = \frac{\hbar^2}{8m}\int \frac{(\vec{\nabla}\rho(\vec{r}))^2}{\rho(\vec{r})}d^3r = \frac{\hbar^2}{2m}\int [\vec{\nabla}\rho^{1/2}(\vec{r})]^2 d^3r, \tag{6}$$

is an important component of the DFT kinetic energy. Indeed, T_W is considered exact in the limit of rapidly varying density ρ [25,26] and nine times large than the second term (the Kirgnitz correction [19]) of the semiclassical expansion (2), which have to be considered as an asymptotic expansion [23].

We present in this letter the universal method to construct the rigorous lower bounds to the T_W which allows us to study a few-fermion system at unitarity.

We rewrite Eq.(6) as

$$T_W[\rho] = \int [\frac{\hbar^2}{2m}[\vec{\nabla}\rho^{1/2}(\vec{r})]^2 + (V(\vec{r},\lambda_1,\lambda_2,...\lambda_M) - V(\vec{r},\lambda_1,\lambda_2,...\lambda_M))]\rho(\vec{r})d^3r, \qquad (7)$$

where $V(\vec{r},\lambda_1,\lambda_2,...\lambda_M)$ is an one-particle potential and λ are parameters.

Introducing an auxiliary Hamiltonian H_{aux} as

$$H_{aux} = -\frac{\hbar^2}{2m}\triangle + V(\vec{r},\lambda_1,\lambda_2,...\lambda_M), \qquad (8)$$

and using

$$< \psi|H_{aux}|\psi > \geq E_G < \psi|\psi >, \qquad (9)$$

we get

$$T_W[\rho] \geq NE_G(\lambda_1,\lambda_2,...\lambda_M) - \int V(\vec{r},\lambda_1,\lambda_2,...\lambda_M)\rho(\vec{r})d^3r, \qquad (10)$$

where E_G is the ground state energy of the auxiliary Hamiltonian (8). Therefore, a set of optimal values of parameters λ_i which yield an optimal value for a lower bound to the $T_W[\rho]$ is given by

$$T_W^{lower}[\rho] = \max_{\lambda_i}[NE_G(\lambda_1,\lambda_2,...\lambda_M) - \int V(\vec{r},\lambda_1,\lambda_2,...\lambda_M)\rho(\vec{r})d^3r] \qquad (11)$$

Now let us collect some lower bounds to the Weizsäcker energy $T_W[\rho]$:

(i) $V(\vec{r},\lambda) = m\lambda^2r^2/2$, $E_G(\lambda) = (3/2)\hbar\lambda$, $\lambda \geq 0$,

$$T_W[\rho] \geq \frac{9}{8}\frac{\hbar^2N^2}{m < r^2 >}. \qquad (12)$$

(ii) $V(\vec{r},\lambda) = -\lambda/r$, $E_G(\lambda) = \lambda^2m/(2\hbar^2)$, $\lambda \geq 0$,

$$T_W[\rho] \geq \frac{\hbar^2}{2mN} < r^{-1} >^2 . \qquad (13)$$

(iii) $V(\vec{r},\lambda_1,\lambda_2,\lambda_3) = (m/2)(\lambda_1^2x^2 + \lambda_2^2y^2 + \lambda_3^2z^2)$, $E_G(\lambda_1,\lambda_2,\lambda_3) = (\hbar/2)(\lambda_1+\lambda_2+\lambda_3)$, $\lambda_i \geq 0$,

$$T_W[\rho] \geq \frac{\hbar^2N^2}{8m}(\frac{1}{< x^2 >} + \frac{1}{< y^2 >} + \frac{1}{< z^2 >}). \qquad (14)$$

(iv) $V(\vec{r},\lambda) = m\lambda^2r^2/2 + \beta(\beta+1)\hbar^2/(2mr^2)$, $E_G = \hbar\lambda(3/2+\beta)$, $\beta \geq 0$, $\lambda \geq 0$,

$$T_W[\rho] \geq \frac{\hbar^2}{2m} < r^{-2} > \frac{3N^2 + < r^2 >< r^{-2} >}{4(< r^2 >< r^{-2} > -N^2)}. \qquad (15)$$

(v) $V(\vec{r},\lambda) = -\lambda/r + \beta(\beta+1)\hbar^2/(2mr^2)$, $E_G = -\lambda^2 m/(2\hbar^2(1+\beta)^2)$, $\beta \geq 0$, $\lambda \geq 0$,

$$T_W[\rho] \geq \frac{\hbar^2}{8m} < r^{-2} > \frac{1}{1 - \frac{<r^{-1}>^2}{N<r^{-2}>})}, \tag{16}$$

where $< r^\alpha >= \int \rho(\vec{r}) r^\alpha d^3r$.

The bounds (12), (13), (15) and (16) where previously proved in Refs. [27,28], while the bound (14) is new.

In Refs.[29,30] and later in Refs.[31-42] the dynamics of strongly interacting dilute Fermi gases (dilute in the sense that the range of interatomic potential is small compared with inter-particle spacing) consisting of a 50-50 mixture of two different states and confined in a harmonic trap $V_{ext}(\vec{r}) = (m/2)(\omega_\perp^2(x^2+y^2) + \omega_z^2 z^2)$ is investigated in the single equation approach to the time-dependent density-functional theory.

Let us come back to the variational formulation of the Kohn-Sham time-dependent theory

$$\delta \int dt < \psi | i\hbar\partial_t - H | \psi >= 0, \qquad 17)$$

where $|\psi>$ is a product of two Slater determinants, one for each internal state built up by the Kohn-Sham orbitals ψ_i, and $H = T + U$ is the LDA Hamiltonian.

Using two approximations

(i) local transform $\psi_i \approx \phi_i \exp(i\hbar\chi/m)$, where ψ_i and χ are real functions,

and

(ii) $< \phi|T|\phi > \approx \frac{\hbar^2}{2m} \int \tau_{TF}(\rho) d^3r + T_W[\rho]$,

where $|\phi>$ is the product of two Slater determinants built on ϕ_i alone, we can derive the DFT equation of Ref.[29]

$$i\hbar\frac{\partial\Psi}{\partial t} = -\frac{\hbar^2}{2m}\nabla^2\Psi + V_{ext}\Psi + V_{xc}\Psi, \tag{18}$$

where $V_{xc}(\vec{r},t) = [\frac{\partial\rho\epsilon(\rho)}{\partial\rho}]_{\rho=\rho(\vec{r},t)}$, $\epsilon(\rho)$ is the ground-state energy per particle of the homogeneous system and $\rho(\vec{r},t) =| \Psi(\vec{r},t) |^2$. In the case of large but finite number of atoms N, at small distances the ratio $| \nabla\rho | / \rho^{4/3}$ is small and both the Kirzhnitz correction and the Weizsäcker correction are negligible. On the contrary, the Weizsäcker correction is expected to determine the asymptotic behavior of the density at large distances. As for the case of relatively small number of atoms, we expect that the Kirzhnitz correction would be a reasonable approximation to the kinetic energy

$$T[\rho] \approx T_{TF}[\rho] + \frac{1}{9} T_W[\rho] d^3r, \tag{19}$$

where $T_{TF}[\rho] = \frac{\hbar^2}{2m}\int(\tau_{TF}(\rho)d^3r$.

Recently, Ref.[41] has considered the following nonlinear equation

$$i\hbar\frac{\partial\Psi_s}{\partial t} = -\frac{\hbar^2}{4m}\nabla^2\Psi + 2V_{ext}\Psi_s + 2V_{xc}\Psi_s, \tag{20}$$

with $|\Psi_s|^2 = \rho/2$.

For the stationary case Eq.(20) reduces to

$$-\frac{\hbar^2}{8m}\nabla^2\Psi + V_{ext}\Psi + V_{xc}\Psi = \mu\Psi \tag{21}$$

which corresponds to the following approximation of the kinetic energy

$$T[\rho] \approx T_{TF}[\rho] + \frac{1}{\kappa}T_W[\rho] \tag{22}$$

with $\kappa = 4$.

For the remainder of this letter we will test the accuracy of the approximation (22) with $\kappa = 9$ and $\kappa = 4$ for few-fermion systems at unitarity in a spherical harmonic trap

$$V_{ext}(\vec{r}) = \frac{m\omega^2 r^2}{2}. \tag{23}$$

In the approximation (22), the ground state energy is given by the minimum of the energy functional

$$J[\Psi] = \frac{1}{\kappa}T_W[\rho] + \int V_{ext}\rho d^3r + \int \epsilon(\rho)\rho d^3r, \tag{24}$$

where $\epsilon(\rho) = (1+\beta)3\hbar^2 k_F^2/(10m)$, $k_F = (3\pi^2\rho)^{1/3}$, $\rho = |\Psi|^2$, and the universal parameter β is estimated to be $\beta = -0.56$ [7].

Introducing an auxiliary Hamiltonian

$$\tilde{H} = 3\frac{\hbar\omega}{2}\sqrt{\frac{\lambda}{\kappa}} + \frac{m\omega^2}{2}(1-\lambda)r^2, \tag{25}$$

we can rewrite Eq.(24) as

$$J[\Psi] = \frac{1}{\kappa}T_W[\rho] + \int(V_{ext} - \tilde{H})\rho d^3r + \int(\tilde{H} + \epsilon(\rho))\rho d^3r.$$

Omission of $(1/\kappa)T_W[\rho] + \int(V_{ext} - \tilde{H})\rho d^3r$ yields our approximation for the ground state energy

$$E \approx \frac{3}{2}N\hbar\omega\sqrt{\frac{\lambda}{\kappa}} + E_{TF}(\sqrt{1-\lambda}\omega), \tag{26}$$

where E_{TF} is the Thomas-Fermi energy which is given by $E_{TF}(\omega) = \sqrt{1+\beta}\hbar\omega(3N)^{4/3}/4$

Projecting $|\Psi>$on the complete basis states $|n>$, obtained from $h|n>= e_n|n>$, where $h = -(\hbar^2/(2\kappa m))\triangle + m\omega^2\lambda r^2/2$, we get

$$< \Psi|h|\Psi >= \sum_n e_n < \Psi|n >< n|\Psi > \geq \frac{3}{2}N\hbar\omega\sqrt{\frac{\lambda}{\kappa}}.$$

Therefore, we conclude that our approximation for energy, given by Eq.(26), is a lower bound to the ground sate energy, Eq.(24). The optimal value of parameter λ which maximizes the energy, Eq.(26), will yield an optimal value of the lower bounds to the ground state energy given by

$$\frac{E}{\hbar\omega} \geq \frac{E^{(-)}}{\hbar\omega} = \frac{3}{2}\frac{N}{\sqrt{\kappa}}\sqrt{1+\frac{(3N)^{2/3}(1+\beta)\kappa}{4}}. \tag{27}$$

For large N, the finite N correction can be written as

$$\frac{E^{(-)}}{\hbar\omega} = \frac{(3N)^{4/3}\sqrt{1+\beta}}{4} + \frac{(3N)^{2/3}}{2\kappa\sqrt{1+\beta}} + ... \tag{28}$$

Bhaduri, Murthy and Brack [43] have recently presented a semiclassical approximation, assuming the particles obey the Haldane-Wu fractional exclusion statistics at unitarity

$$\frac{E^{BMB}}{\hbar\omega} = g^{1/3}\frac{(3N)^{4/3}}{4} + g^{-1/3}\frac{(3N)^{2/3}}{8} + ..., \tag{29}$$

where the statistical parameter g is related to the universal parameter β by the relation $g = (1+\beta)^{3/2}$. We note here that for the case of $\kappa = 4$ [41] our lower bound, Eq.(28), agrees with the expression of Ref.[43].

To calculate upper bounds, $E^{(+)}$, we employing Fetter's trial functions [44]

$$\rho^{1/2}(\vec{r}) = c[1-(1-q)(\gamma r)^2]^{\frac{1}{1-q}}, \tag{30}$$

where γ, q are the variational parameters and c is the normalization constant, to minimize the functional J, Eq.(24).

From Table I, we can see that $E^{(-)} > E^{MC}$ for $\kappa = 4$ and $2 \leq N/2 \leq 14$. Therefore

$$\frac{|E(\kappa=4) - E^{MC}|}{E^{MC}} > \triangle(\kappa = 4),$$

where $\triangle(\kappa = 4) = (E^{(-)}(\kappa = 4) - E^{MC})/E^{MC}$ and $E(\kappa)$ is the exact ground state solution of Eq.(24). For $\kappa = 9$ and $3 \leq N/2 \leq 14$, $E^{(+)} \leq E^{MC}$ (see Table I), that is why

$$\frac{|E^{MC} - E(\kappa=9)|}{E^{MC}} > \triangle(\kappa = 9),$$

where $\triangle(\kappa = 9) = (E^{MC} - E^{(+)}(\kappa = 9))/E^{MC}$.

We can, therefore, state that $\triangle(\kappa)$ is the rigorous lower bound to the accuracy of the approximation (22) with $\kappa = 4$, $2 \leq N/2 \leq 14$ and $\kappa = 9$, $3 \leq N/2 \leq 14$. As for the lower $E^{(-)}$ and the upper $E^{(+)}$ bounds, they provide the actual solution of Eq.(24), $(E^{(+)} + E^{(-)}/2$ within $\pm\delta$ accuracy, with $\delta < 1\%$ for both $\kappa = 9$ and $\kappa = 4$ and for $2 \leq N/2 \leq 15$.

The predictions of Eq.(24) for the ground state energy and results of Ref.[10] are shown in Fig.1. A very good agreement between the Kirgnitz approximation, $\kappa = 9$, and the *ab initio* calculations of Ref.[10] can be seen for $2 \leq N/2 \leq 4$. However, for $12 \leq N/2 \leq 15$ the $\kappa = 4$ approximation gives better results than the $\kappa = 9$ approximation.

Finally, we note that the kinetic energy functionals assumed here are not unique. In our future work we will consider other possible forms.

In summary, we have constructed the rigorous lower bounds to the Weizsäcker energy. As example of application, we have studied few-fermion systems at unitarity consisting of 50-50 mixture of two different states and confined in a spherical harmonic trap. The rigorous lower bounds to the accuracy of the method are derived. We have tested the kinetic energy functionals by comparisons with *ab initio* calculations and have found that while the second order gradient expansion is a very accurate for relatively small N, Fig.1 indicates that the $\kappa = 4$ approximation provides significantly better results for larger N.

Acknowledgments

I thank N.J. Giordano and W.L. Fornes for providing the opportunity to finish this work.

Table I. The energies $E^{(-)}$, $E^{(+)}$ and the energy calculated within the fixed-node diffusion Monte Carlo method, E^{MC} [10], all in units of $\hbar\omega$ for $N \leq 30$ [see the text for further details].

$N/2$	$E^{(-)}(\kappa=4)$	$E^{(+)}(\kappa=4)$	E^{MC}	$E^{(-)}(\kappa=9)$	$E^{(+)}(\kappa=9)$	$\triangle(\kappa=4)$	$\triangle(\kappa=9)$
2	5.455	5.561	5.05	4.976	5.062	8.0%	-
3	9.025	9.198	8.64	8.378	8.513	4.5%	1.5%
4	12.954	13.196	12.58	12.157	12.341	3.0%	1.9%
5	17.182	17.495	16.81	16.247	16.479	2.2%	2.0%
6	21.670	22.055	21.28	20.605	20.885	1.8%	1.9%
7	26.390	26.846	25.92	25.202	25.530	1.8%	1.5%
8	31.318	31.848	30.88	30.014	30.389	1.4%	1.6%
9	36.440	37.043	35.97	35.024	35.445	1.3%	1.5%
10	41.741	42.416	41.30	40.216	40.684	1.1%	1.5%
11	47.208	47.956	46.89	45.578	46.093	0.7%	1.7%
12	52.833	53.653	52.62	51.100	51.662	0.4%	1.8%
13	58.605	59.499	58.55	56.775	57.381	0.1%	2.0%
14	64.519	65.485	64.39	62.592	63.244	0.2%	1.8%
15	70.567	71.606	70.93	68.546	69.243	-	2.4%



Fig.1. Ground state energy per particle of few-fermion systems at unitarity in a spherical harmonic trap in units of $\hbar\omega$ as a function of number of atoms N. The dashed and the dotted-dashed lines represent $(E^{(-)} + (E^{(+)})/2$ for $\kappa = 9$ and $\kappa = 4$, respectively. The circular dots indicate results of Ref.[10].

References

[1] O'Hara K M, Hemmer S L, Gehm M E, S R Granade S R, and Thomas J E 2002 *Science* **298**,2179

[2] Bartenstein M, Altmeyer A, Riedl S, Jochim S, Chin C, Denschlag J H, and Grimm R 2004 *Phys. Rev. Lett.* **92**, 120401

[3] Bourdel T, Khaykovich L, Cubizolles J, Zhang J, Chevy F, Teichmann M, Tarruell L, Kokkelmans S J J M F, and Salomon C 2004 *Phys. Rev. Lett.* **93**, 050401

[4] Bishop R F 2001 *Int. J. Mod. Phys.* **15**, iii

[5] Astrakharchik G E, Boronat J, Casulleras J, and Giorgini S 2004 *Phys. Rev. Lett.* **93**, 200404

[6] Chang S Y and Pandharipande V R 2005 *Phys. Rev. Lett.* **95**, 080402

[7] Carlson J, Chang S-Y, Pandharipande V R, and Schmidt K E 2003 *Phys. Rev. Lett.* **91**, 050401

[8] Carlson J and Reddy S 2005 *Phys. Rev. Lett.* **95**, 060401

[9] Chang S Y, and Bertsch G F 2007 *Phys. Rev.* A**76**, 021603(R)

[10] Blume D, von Stecher J, and Greene C H 2007 *Phys. Rev. Lett.* **99**, 233201

[11] von Stecher J, Greene C H, and Blume D 2007 *Phys. Rev.* A**76**, 053613

[12] Blume D 2008 *Phys. Rev.* A**78**, 013635

[13] von Stecher J, Greene C H, and Blume D 2008 *Phys. Rev.* A**77**, 043619

[14] Kohn W, and Sham L J 1965 *Phys. Rev.* **140**, A1133

[15] Hohenberg P, and Kohn W 1964 *Phys. Rev.* **136**, B1133

[16] Yu Y, and Bulgac A 2003 *Phys. Rev. Lett.* **90**, 222501

[17] Bulgac A 2002 *Phys. Rev.* C**65**, 051305(R)

[18] Bulgac A 2007 *Phys. Rev.* A**76**, 040502(R)

[19] Kirgnitz D A 1967 *Field Theoretical Methods in Many Body Systems* (London: Pergamon Press)

[20] Brack M, and Bhaduri R K 1997 *Semiclassical Physics* (Reading MA: Addison-Wesley)

[21] Brack M, Jennings B K, and Chu P H 1976 *Phys. Lett.* B**65**, 1

[22] Hodges C H 1973 *Can. J. Phys* **51**, 1428

[23] Guet C, and Brack M 1980 *Z. Phys.* A**297**, 247

[24] von Weizsäcker C F 1935 *Z. Phys.* **96**, 431

[25] Parr R G, and Yang W 1989 *Density-Functional Theory of Atoms and Molecules* (New York: Oxford University Press)

[26] Dreizler R M, and Gross E K U 1990 *Density Functional Theory: An Approach to the Quantum Many-Body Problem* (Berlin: Springer-Verlag

[27] Gadre S R and Pathak R K 1982 *Phys. Rev.* A**25**, 668

[28] Romera E and Dehesa J S 1994 *Phys. Rev.* A**50**, 256

[29] Kim Y E and Zubarev A L 2004 *Phys. Rev.* A**70**, 033612

[30] Kim Y E and Zubarev A L 2004 *Phys. Lett.* A**327**, 397

[31] Kim Y E and Zubarev A L 2005 *Phys. Rev.* A**72**, 011603(R)

[32] Kim Y E and Zubarev A L 2005 *J. Phys. B: At. Mol. Opt. Phys.* **38**, L243

[33] Manini N and Salasnich L 2005 *Phys. Rev.* A**71**, 033625

[34] Ghosh T K and Machida K 2006 *Phys. Rev.* A**73**, 013613

[35] Diana G, Manini N, and Salasnich L 2006 *Phys. Rev.* A**73**, 065601

[36] Yin J and Ma Y-L 2006 *Phys. Rev.* A**74**, 013609

[37] Salasnich L and Manini N 2007 *Laser Phys.* **17**, 169

[38] Zhou Y and Huang G 2007 *Phys. Rev.* A**75**, 023611

[39] Ma Y-L and Huang G 2007 *Phys. Rev.* A**75**, 063629

[40] Wen W and Huang G 2007 *Phys. Lett.* A**362**, 331

[41] Wen W, Zhou Y and Huang G 2008 *Phys. Rev.* A**77**, 033623

[42] Adhikari S K 2008 *Phys. Rev.* A**77**, 045602

[43] Bhaduri R K, Murthy M V N and Brack M 2008 *J. Phys. B: At. Mol. Opt. Phys.* **41**, 115301

[44] Fetter A.L. 1997 *J. Low. Temp. Phys.* **106**, 643